

Purina

THE SMALL FLOCK PROGRAM




Broiler/Roaster



Turkeys



Layers



Purina Start & Grow Complete and balanced feed for chicks and growing pullets. It will give your birds a healthy start and vigorous growth from day one.

Purina Growena A complete and balanced feed suitable for growing broiler chickens, pullets, turkeys and ducks

Purina Meat Builder A complete, 18% crude protein ration formulated for broiler and roaster chickens. This is a high performance feed for quick growth of meat-type birds (chickens and turkeys).

Purina Layena This high energy ration is formulated with extra vitamins and minerals for top egg production and shell quality in egg laying hens, ducks and geese.

Feeding Program For Broilers/Roasters

Chicks should be fed on a free-choice basis and should have fresh feed at all times. After 5 weeks of age, double the feeder space and water capacity. NOTE: Broilers are usually processed at about 2 kg (4.4 lbs) while roasters can be processed at 3.5 kg (8 lbs). Pullets are usually smaller and lighter than cockerels of the same age.

Feeding Program For Turkeys

Provide one tube feeder for every 10-12 turkeys. If trough feeders are used, 20 cm (8 in.) are required per bird. Turkey poults must have easy access to feed during the first week of life. Try using egg cartons, since they make excellent feeders for starting turkeys. Clean fresh water should be available for the birds on a free-choice basis at all times. At 6 weeks of age, increase the floor space up to 0.5 sq. m (5 sq. ft.) per bird and double the feeder space and water capacity. NOTE: Turkeys will achieve a weight of about 5 kg (11 lbs) at about 14 weeks but toms can be grown to more than 13 kg (29 lbs) in 25 weeks. Hens are generally smaller at the same age.

Feeding Program For Egg Laying Birds

Birds should be fed on a free-choice basis at all times and fresh, room-temperature water should always be available. After 20 weeks, feeding and watering space should be increased to one tube feeder for every 10-15 hens or 10 cm (4 in) feeder space per hen. There should be one nest for every 5-7 hens. Supplemental light should be provided. Ensure that the hens receive 14 -17

hours of continuous light per day to stay in egg production. If the lights are turned on at 6 am, most hens will lay their eggs by 11 am. Supplemental heat should be provided when the temperature is less than 15°C (60°F) to maintain egg production.

Feeding Program For Ducks and Geese

Fresh, clean water should be available for birds at all times. Provide one tube feeder for every 25 ducks and one for every 10-12 geese. If trough feeders are used, 5 cm (2 in) are required by ducks and 20 cm (8 in) by geese.

GUARANTEED ANALYSIS

Products	Protein (%)	Fat (%)	Fibre (%)	Na (%)	Ca (%)	P (%)
6039	20	4.0	4.5	0.18	1.00	0.75
6032	15	2.0	6.0	0.16	0.85	0.70
6054	18	3.0	5.0	0.16	0.90	0.70
6055	18	3.0	5.0	0.16	0.90	0.70
6498	17	1.5	4.5	0.16	3.80	0.60
8016	24	3.0	6.0	0.15	1.30	0.90

For more information, contact your local Purina Dealer, call 1-800-567-CHOW(2469), email info@agribands.ca or visit our website www.agripurina.ca



Purina